

PROGRAM PRACTICES SUBMISSION FORM

IMMUNIZATION PROGRAM PRACTICES INFORMATION

Name: Melissa K. Fankhauser, MPH/ Kelly Moore, MD, MPH Email address: <u>melissa.fankhauser@tn.gov</u>; <u>Kelly.moore@tn.gov</u> Program: Tennessee Immunization Program (TIP)

Does AIM have permission to share this information on the publicly accessible AIM website? All materials submitted will be posted on the AIM website. Please note, the
manuscript for this project is in press; we request that materials be posted after
publication.

BULL'S-EYE AWARD

The **Bulls-Eye Award for Innovation and Excellence in Immunization** recognizes immunization strategies that "hit their mark" and achieve immunization goals with special consideration for practices that are innovative and easily replicated. Each year, AIM awards three programs the Bull's-Eye Award. Only those practices with activities that are **currently ongoing** or **concluded during 2016** (including implementation, follow-up, and/or evaluation activities) will be considered for the award.

Would you like for this submission to be considered for the 2017 AIM Bull's- Eye	<u>X</u> Yes	No
Award?		

PROGRAM PRACTICE INFORMATION

Title

The Tennessee Immunization Program 3-Star Report: Using Audit and Feedback to Reduce Missed Opportunities for Preteen Immunization in Local Public Health Clinics

Keywords (up to 5 main terms/phrases that describe the practice) Adolescent immunization, HPV vaccine, public health, information systems, quality improvement

Is this practice Evidence / Guideline Based? (if yes, include reference below) _X_Yes __No

Reference:

1. President's Cancer Panel. Accelerating HPV Vaccine Uptake: Urgency for Action to Prevent Cancer. President's Cancer Panel Annual Report 2012–2013. 2014. Available at: <u>http://deainfo.nci.nih.gov/advisory/pcp/annualReports/</u>HPV/index.htm. Published 2014.

2. Jacob V, Chattopadhyay SK, Hopkins DP, et al. Increasing coverage of appropriate vaccinations: a community guide systematic economic review. Am J Prev Med. 2016;50(6):797–808.

3. Briss PA, Rodewald LE, Hinman AR, et al. Reviews of evidence regarding interventions to improve vaccination coverage in children, adolescents, and adults. The Task Force on Community Preventive Services. Am J Prev Med.

2000;18(suppl 1):97-140.

Background (scope of the immunization need or problem)

In the CDC's 2014 National Immunization Survey-Teen, Tennessee performed poorly in the measures of HPV vaccination for both males and females aged 13–17 years. Among males, Tennessee fell in the bottom quartile of states for both initiation (30.5%, \pm 8.5) and series completion (14.0%, \pm 6.6); among females, the point estimate of HPV initiation (47.8%, \pm 9.8) and series completion (20.1%, \pm 6.7) was among the lowest in the nation. Tdap and Men-ACWY coverage was 86.0% (\pm 4.5) and 74.0% (\pm 5.8), respectively.

Very few children receive all immunizations at a local health department (LHD) in Tennessee; immunizations are provided opportunistically. The state Tdap requirement for 7th grade entry provides the ideal opportunity to administer all preteen vaccines at one visit ("bundling").

Program Practice Description

Describe the practice goals and objectives.

In 2013, TIP developed the monthly Vaccines for Children (VFC) Preteen Public Health Vaccination Report, abbreviated "3-Star Report", as a continuous quality improvement (CQI) measure for local health departments (LHDs) across the state aimed at increasing bundled administration of Tdap, Men-ACWY, and HPV vaccines during the same visit, which has been shown to improve on-time completion of vaccines. The 3-Star Report evaluates preteen vaccines given at immunization visits of VFC-eligible children aged 11 through 13 years who received a Tdap, a MenACWY, and/or an HPV vaccine at an LHD. Visits are categorized as 1-Star (one vaccine given), 2-Star (two vaccines given), or 3-Star (all three given). This simple, low-cost program leverages the existing information system used by the LHDs. Because this system allows only evaluation of the encounter and not the child's full immunization history, performance targets were not established. The objective is to increase awareness, motivation and performance among LHD nurses at each encounter by measuring and reporting 3-Star performance at the county, regional and state level every month.

What were the main implementation activities?

LHDs in Tennessee use the Patient Tracking and Billing Management Information System (PTBMIS) to collect data on immunization visits; data are extracted from this system by the TIP epidemiologist each month. The epidemiologist designed the report and programmed production and distribution of the report in SAS version 9.3. Each month, the TIP epidemiologist runs the report and it is distributed to relevant regional and state public health leadership. TIP does not dictate how the report is used or shared: LHD use varies.

Where and when did the practice take place?

The report was first distributed in April 2013 and the activity is ongoing.

How much staff time was involved?

Creating the report took the TIP epidemiologist approximately 2 months while working on other job duties. It takes approximately 30 minutes to generate and distribute the 3-Star report each month.

What were the costs associated with the activity? What was the funding source?

Federal Section 317 funds are used to pay for the epidemiologist and SAS license: no funding was specifically requested to create the report. The only cost of the ongoing report is the 30 minutes of staff time per month.

Identify the target population that the practice affected.

The report is distributed to all 13 regional and metropolitan HDs for further dispersal to local clinic staff. It is intended to impact the performance of nurses at the LHD clinic level. The children targeted are those 11 through 13 eligible for VFC who come to an LHD for an immunization visit. Between September 2015 and August 2016, 16,156 eligible children visited LHDs, with 8,431 (52%) visiting in July and August, when parents of 7th grade students rush to provide schools with state-required proof of Tdap immunization.

If partners were involved, include who was involved, and how.

Regional and local HDs use the report in a variety of ways. Some have leadership teams that track performance, aiming to have the highest state 3-Star rates each month. One introduced a temporary incentive program, giving a gift card and certificate to the nurse who gave the most 3-Star visits in a month

(resulting in a rise in 3-star visits in that region from 7 percent to 50 percent over 2 months).

Timeframe of Implementation (Start and Stop Dates)

The 3-Star report has been produced each month since April 2013.

Evaluation Data: Was the implementation and/or effectiveness of this	<u>X</u> Yes	No	Limited
practice assessed? (if "yes" or "limited," provide any data that is available)			

Data: See charts below. Figure 3 (left) of the in-press manuscript shows the year-over-year increases in 3-Star visits each summer during the back to school rush. The 3-Star rates went up by 63.9% between July 2013 (25.5%) and July 2016 (41.7%). Figure 4 (right) shows that the proportion of non-HPV 1-Star visits (typically Tdap-only visits) have declined compared to the same month in the previous year since introduction of the report. Although results represent correlation and not causation, bundling in LHDs has clearly improved.



Figure 3. Percentage of 3-Star visits each month among VFC-eligible children aged 11–13 years who received a Tdap, Men-ACWY, and/or HPV vaccine at a Tennessee LHD, January 2013 through August 2016. Months of the year are on the x-axis, and percentages are on the y-axis.

Figure 4. Percentage of non-HPV 1-Star visits each month among VFCeligible children aged 11–13 years who received a Tdap, Men-ACWY, and/ or HPV vaccine at a Tennessee LHD, January 2013 through August 2016. Months of the year are on the x-axis, and percentages are on the y-axis.

Conclusions / Lessons Learned / Key Factors for Success

The 3-Star Report is easy to maintain and has been correlated with year-over-year increases in 3-Star visits and the corresponding decreases in non-HPV 1-Star visits each summer during the back to school rush. The report has generated a healthy sense of competition among regions, some of which now achieve 3-Star visit rates of at least 50% consistently.

This CQI measure could be used by other programs or individual clinics that collect data on immunization visits of preteens. Should programming not be available, the measure is not complicated and it is feasible for a clinic to track its own 3-Star visit rates on paper.

Check if any of the following are being submitted to complement your submission:

(All materials will be posted on the AIM website)

*A manuscript is now in press. It will be published soon in a special supplement of the open source, peer-reviewed journal, *Biomedical Informatics Insights*. As soon as it is published, we will provide the link to the full manuscript. **Email the completed form to Mary Waterman (<u>mwaterman@immunizationmanagers.org</u>) with the subject line "Program Practices Submission."

		1-Star				2-Star		3-Star	
County	Total Number *	Number Non-HPV	% Non-HPV	Number HPV	% HPV	Number	%	Number	%
Benton	6	0	0.0%	6	100.0%	0	0.0%	0	0.0%
Carroll	12	2	16.7%	5	41.7%	2	16.7%	3	25.0%
Chester	4	0	0.0%	3	75.0%	0	0.0%	1	25.0%
Crockett	15	0	0.0%	13	86.7%	0	0.0%	2	13.3%
Decatur	6	0	0.0%	3	50.0%	1	16.7%	2	33.3%
Dyer	7	2	28.6%	2	28.6%	0	0.0%	3	42.9%
Fayette	11	1	9.1%	7	63.6%	1	9.1%	2	18.2%
Gibson	62	7	11.3%	14	22.6%	13	21.0%	28	45.2%
Hardeman	13	2	15.4%	6	46.2%	0	0.0%	5	38.5%
Hardin	18	1	5.6%	16	88.9%	0	0.0%	1	5.6%
Haywood	17	0	0.0%	14	82.4%	0	0.0%	3	17.6%
Henderson	6	0	0.0%	6	100.0%	0	0.0%	0	0.0%
Henry	3	1	33.3%	1	33.3%	1	33.3%	0	0.0%
Lake	1	0	0.0%	1	100.0%	0	0.0%	0	0.0%
Lauderdale	28	14	50.0%	3	10.7%	4	14.3%	7	25.0%
McNairy	25	6	24.0%	11	44.0%	5	20.0%	3	12.0%
Obion	9	0	0.0%	8	88.9%	0	0.0%	1	11.1%
Tipton	19	3	15.8%	7	36.8%	3	15.8%	6	31.6%
Weakley	5	0	0.0%	5	100.0%	0	0.0%	0	0.0%
	267	39	14.6%	131	49.1%	30	11.2%	67	25.1%

3-Star Report for October 2016, Counties in WTR

* Total number of fully VFC-eligible children (excluding under-insured) aged 11 through 13 years who received at least one of the following 3 vaccines: Tdap, MCV4, and/or HPV.



1-Star			2-Star		3-Star				
	Total	Number	%	Number	%				
Region	Number *	Non-HPV	Non-HPV	HPV	HPV	Number	%	Number	%
Statewide	1272	175	13.8%	674	53.0%	167	13.1%	256	20.1%

3-Star Report for October 2016, Statewide

* Total number of fully VFC-eligible children (excluding under-insured) aged 11 through 13 years

who received at least one of the following 3 vaccines: Tdap, MCV4, and/or HPV.





Tennessee's 3-Star Report: Using Available Data Systems to Reduce Missed Opportunities to Vaccinate Preteens



Kelly L. Moore¹, Melissa K. Fankhauser¹ and Pamela C. Hull²

¹Tennessee Department of Health, Immunization Program, Nashville, TN, USA. ²Department of Medicine, Division of Epidemiology, Vanderbilt University Medical School, Nashville, TN, USA.

Supplementary Issue: Use of Biomedical Informatics for Improving Vaccine Uptake and Adherence

ABSTRACT: All preteens should receive tetanus-diphtheria-pertussis vaccine (Tdap), quadrivalent meningococcal vaccine (Men-ACWY), and the human papillomavirus (HPV) cancer vaccine series. In Tennessee, HPV vaccination rates have stagnated at low levels for a decade. Three fundamental strategies to reduce missed opportunities for immunization include administering all recommended vaccines at the same visit, making strong recommendations for vaccines, and auditing and feedback. In Tennessee, during each summer, a surge of preteens visit local health departments (LHDs) to receive a required Tdap vaccine before entering seventh grade, presenting an opportunity to administer Men-ACWY and HPV. The Tennessee Immunization Program (TIP) coined the term "3-Star visit" for such encounters and developed a monthly report to track them using data from the Patient Tracking Billing Management Information System (PTBMIS) used by LHDs across Tennessee. Implementation of this quality improvement report has correlated with a substantial increase in 3-Star visits from 2013 to 2016, particularly during the summer months.

KEYWORDS: adolescent immunization, HPV vaccine, public health, information systems, quality improvement

SUPPLEMENT: Use of Biomedical Informatics for Improving Vaccine Uptake and Adherence

CITATION: Moore et al. Tennessee's 3-Star Report: Using Available Data Systems to Reduce Missed Opportunities to Vaccinate Preteens. *Biomedical Informatics Insights* 2016:8(S2) 15–21 doi: 10.4137/BII.S40207.

TYPE: Technical Advance

RECEIVED: August 2, 2016. RESUBMITTED: November 03, 2016. ACCEPTED FOR PUBLICATION: November 05, 2016.

ACADEMIC EDITOR: John P. Pestian, Editor in Chief

PEER REVIEW: Seven peer reviewers contributed to the peer review report. Reviewers' reports totaled 1024 words, excluding any confidential comments to the academic editor.

FUNDING: PCH received support from National Institutes of Health, grant numbers U54 CA163072 and P30 CA068485-18S3. The authors confirm that the funder had no influence over the study design, content of the article, or selection of this journal.

COMPETING INTERESTS: KLM discloses: salary funding via grants to the Tennessee Department of Health from the US CDC; membership on the CDC Advisory Committee on Immunization Practices, including travel expenses and stipend; honorarium and travel

Introduction

Vaccines recommended by the U.S. Centers for Disease Control and Prevention for preteens and teens can prevent serious and life-threatening conditions¹; however, immunization providers have failed over the last decade to reach the high rates of immunization in this age group that have been achieved for vaccines given in infancy.^{2,3} Three vaccines are recommended for all children aged 11–12 years: a booster dose of tetanus– diphtheria–pertussis (Tdap) vaccine,⁴ a dose of quadrivalent meningococcal vaccine (Men-ACWY),⁵ and a three-dose series of vaccine against select strains of the human papillomavirus (HPV) capable of causing a spectrum of diseases, including several types of cancer in men and women.^{6–9}

Despite the burden of preventable disease, preteen immunization coverage for HPV has, in particular, remained unacceptably low, even as Tdap and Men-ACWY coverage rates have increased.² In the CDC's 2014 National Immunization Survey-Teen, Tennessee performed poorly in the measures of HPV vaccination for both males and females aged 13–17 years. Among males, Tennessee fell in the bottom quartile of states for both initiation (30.5%, ±8.5) and series completion expenses from the National Foundation for Infections Diseases; travel expenses from the Association of Immunization Managers; and travel expenses from the Tennessee Academy of Family Physicians. MKF discloses salary funding via grants to the Tennessee Department of Health from the US CDC. PCH disclose no potential conflicts of interest.

CORRESPONDENCE: Kelly.moore@tn.gov

 $\label{eq:copyright: the output} \begin{array}{l} \mbox{COPYRIGHT: } \textcircled{\sc b} \mbox{the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License. \end{array}$

Paper subject to independent expert blind peer review. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to antiplagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Published by Libertas Academica. Learn more about this journal.

(14.0%, ± 6.6); among females, the point estimate of HPV initiation (47.8%, ± 9.8) ranked 49th among the 50 states and series completion (20.1%, ± 6.7) was 50th of 50 states, while Tdap and Men-ACWY coverage was 86.0% (± 4.5) and 74.0% (± 5.8), respectively, which is close to national averages.²

Factors contributing to low HPV vaccine uptake include misinformation raising scientifically unsubstantiated public concerns about its safety, an erroneous belief that the vaccine could lead to early sexual activity or is not needed until sexual activity is expected, and the lack of a strong immunization provider recommendation for administering the vaccine at the age of 11 or 12 years, as recommended by experts.^{10–13} In the face of an increasing number of young people missing out on this important development in cancer prevention, numerous strategies have been recommended to improve timely HPV vaccination, with several outlined by the President's Cancer Panel Annual Report 2012-2013.14 Included among these is an emphasis on giving HPV vaccine when other vaccines are given, a version of which is the principle of "bundling," or administering all three routine adolescent vaccines at the same visit.¹⁴⁻¹⁶ By giving all age-appropriate vaccines on the same day and recommending them in the same way, as in the infant immunization schedule, clinic staff can avoid implying that HPV vaccine is somehow optional, questionable, or exceptional. Another principle is the importance of a strong provider recommendation; most parents will follow their healthcare provider's advice regarding immunizations if the recommendation is clear and unequivocal.^{14,17} In addition, immunization provider audit and feedback is a well-established, evidencebased, and cost-effective strategy for increasing immunization coverage in general, consisting of periodically assessing providers' retrospective performance and sharing this information with them.^{18,19}

The Tennessee Immunization Program (TIP) of the Tennessee Department of Health (TDH) developed a quality improvement (QI) program for local health departments (LHDs) across the state aimed at increasing administration of Tdap, Men-ACWY, and HPV vaccines during the same visit at the recommended ages of 11 and 12 years. This simple, low-cost QI program uses audit and feedback focused on bundling of adolescent vaccines, leveraging the existing information system used by the LHDs. This article describes the development, implementation, and outcomes of the QI program as a public health practice case study. Other states or primary care practices could adapt this simple approach to their existing information systems to increase the administration of bundled adolescent vaccines.

Methods

Overview of QI program. Across Tennessee, LHDs offer all recommended childhood immunizations, including adolescent vaccines, through the Vaccines for Children (VFC) federal entitlement program, eliminating cost as a barrier to access. Although few Tennessee children receive all immunizations at an LHD,²⁰ the LHDs are a convenient option for parents during the back-to-school rush, when all students entering seventh grade in Tennessee, who are typically around 12 years old, must have received a Tdap booster vaccine before classes begin in August.²¹ The rush creates an opportunity for LHD public health nurses to administer all three recommended adolescent vaccines at once. Training around the importance of evidence-based principles of bundling and strong recommendations are provided annually to regional THD immunization representatives, who serve as both auditors and coaches to the VFC clinics in their regions. In addition, all VFC Program participants review strategies for increasing immunization during an annual immunization review conference conducted by state and CDC immunization program staff. To help them remember the three recommended vaccines, the term "3-Star" visit, a reference to the three stars on the state flag, was coined by the TIP in early 2013 to describe a complete first preteen immunization visit.

From the concept of the 3-Star visit came a desire to track these visits to promote performance improvement through regular feedback. In April 2013, the TIP developed



a succinct monthly report to track 1-, 2-, and 3-Star visits by leveraging the standard statewide data available on immunizations given at any LHD in the Patient Tracking Billing Management Information System (PTBMIS). Since then, TIP has distributed the monthly report with feedback to state and regional clinical leadership on the performance of their LHDs, tracking trends over time and comparing regions to one another to support their internal efforts to improve and create healthy competition.

Reports are distributed electronically each month to recipients that include state Community Health Services and Family Health and Wellness Division leadership, to key state leaders interested in HPV vaccine promotion, and to regional public health nursing and medical leadership. Recipients use and share the information with LHD clinic staff as desired to evaluate and improve public health practice: no specific response is required by the program.

Population. The monthly 3-Star Report includes children eligible for the VFC Program aged 11-13 years who presented during the month to any LHD and received at least one routine preteen vaccine (Tdap, HPV, or Men-ACWY) during that encounter. Because the report's intent is to reflect as purely as possible public health nurse performance, children for whom out-of-pocket costs at the LHD could be a barrier, including commercially insured children, were excluded from the assessment. Children under the age of 19 years are entitled to VFC vaccines if they are uninsured, Medicaid eligible, American Indian/Alaskan Native, or if their insurance plan does not cover vaccination as a benefit. Roughly half of all preteens are eligible for VFC vaccines. Underinsured VFCeligible children were excluded from 3-Star reports because they would be VFC-eligible only for a vaccine not covered by insurance.

Information system. All LHDs in Tennessee, until recently, have used PTBMIS for patient and service tracking. PTBMIS is COBOL based and uses a DB2 database manager that runs on i/OS for an iSeries AS400 computer from International Business Machines (IBM®). PTBMIS is a highly networked and distributed environment involving 13 regional computers into which regional public health data are entered and a 14th computer at the state level that integrates these 13 separate systems for statewide reporting. This system was first installed in 1987 and has been used in its present form statewide since 1998. In 2014, one of the 13 regions, a single metropolitan county health department, replaced PTBMIS with a new electronic health record (EHR) system, resulting in its absence from original 3-Star reports from May through September 2014. As a result of county clinic staff demand, this LHD began exporting the necessary data from their EHR to a Microsoft Excel file and securely emailing the file to TIP. Since then, the county data have been appended to the iSeries data from the remaining 12 regional PTBMIS systems for analysis. Summary data presented here include this county's data for the entire reporting period.

The report. An automated process for data analysis, report generation, and e-mail distribution was developed using SAS® version 9.3 (SAS Institute Inc.). Each month, as shown in Figure 1, data are extracted and analyzed on every immunization visit by an eligible child during the previous month. Variables include clinic location, appointment date, patient date of birth, VFC eligibility status, and administered immunizations. The generic vaccine CVX codes included in the analysis are 115 for Tdap, 114 and 136 for Men-ACWY, and 62, 118, and 165 for HPV. The number and proportion of visits that were 1-Star (one of the three vaccines given), 2-Star (two of the three vaccines given), and 3-Star (all three



vaccines given) are obtained at the county, regional, and state levels. Results are presented in tabular and graphic forms. Reports are output in portable document format (PDF) and distributed via e-mail to state and regional users. A total of 14 reports are generated: one for each public health region and one statewide report. To ensure that recipients understand the report, a brief explanatory document accompanies it each time it is distributed.

This article presents a sample 3-Star Report for one public health region report for illustration purposes in Figure 2. In addition, descriptive trends in the monthly proportion of 3-Star Visits and Non-HPV 1-Star Visits are reported for January 2013 to August 2016. The proportion of visits was used rather than the absolute number of visits to adjust for differences in population size and patient volume across regions and across time. The numerators and denominators that comprise the proportions are provided in the Supplementary tables. Since this was a QI program in the context of public health practice and not a hypothesis-driven research study with a controlled research design, statistical tests of changes in trends were not performed. In a practice context, the observed changes over time were of programmatic significance.

Results

Figure 2 illustrates the 3-Star Report layout using data from the counties of the Department of Health Northeast region (NER) in August 2016 as an example. The report begins with a county-level data table of all eligible immunization visits by children 11-13 years of age where at least one of the three vaccines was administered and the proportion of those visits that were 1-, 2-, or 3-Star visits; a total of 269 eligible children in this region were evaluated. One-Star visits are broken down by HPV-only visits and other 1-Star visits. The chart at the bottom of the first page graphically represents the region's results for the current month and the previous 12 months. Bars represent the number of non-HPV 1-Star (red), HPV-only (green/red hatched), 2-Star (yellow), and 3-Star (green) visits each month. The left y-axis represents the scale of numbers of patients. Stars connected by a line represent the percentage of 3-Star visits each month. The right y-axis is the scale of the 3-Star percentage line.

The second page of the sample report provides statewide information for the month. A statewide data table at the top provides raw numbers and proportions of each type of eligible visit: 5,192 eligible children were included in the statewide report in August 2016. The bar chart showing the results for each of the 13 public health regions is shown separately from the percent of 3-Star visits in each region to simplify interpretation. A straight line indicates that 42% (2,179 of 5,175) of all assessed visits statewide were 3-Star visits.

Figure 3 displays statewide results for the monthly proportion of 3-Star visits (solid lines) for 2013–2015 and for the first six months of 2016. The initial report of April 2013 results was distributed in May 2013; due to PTBMIS changes





3-star report for August 2016, Statewide



Figure 2. Example of the 3-Star Report, August 2016, Tennessee Department of Health Northeast Region (NER). Page 1 (left image): a table of results for each county within NER sits above a chart showing the regional results of the 268 eligible patients for the current month and previous 12 months (*x*-axis). In the chart, bar height corresponds to the number of eligible patients (left *y*-axis). Colors within each bar represent the proportion of each type of visit: non-HPV 1-Star (red); HPV-only (green/red hatched); 2-Star (yellow); 3-Star (green). The black line with stars tracks the percentage of 3-Star visits over time (right *y*-axis). Page 2 (right image) displays the statewide results table for the 5,175 eligible patients seen in August 2016; the upper chart shows the number of eligible visits that month in each of the 13 public health regions, using the same colors as the first chart. The bottom chart shows the individual percentages of 3-Star visits in each region for the month, with a horizontal line representing the statewide result of 42% 3-Star visits in August 2016.

to vaccine coding in October 2012, it was not possible to obtain comparable baseline data for the summer of 2012. The year-over-year increases in 3-Star visits each summer during the back-to-school rush are clear, with a 63.9% increase in



Figure 3. Percentage of 3-Star visits each month among VFC-eligible children aged 11–13 years who received a Tdap, Men-ACWY, and/or HPV vaccine at a Tennessee LHD, January 2013 through August 2016. Months of the year are on the *x*-axis, and percentages are on the *y*-axis.

the proportion of 3-Star visits during the peak month of July between 2013 (978 of 3,831 eligible children, or 25.5%) and 2016 (1,352 of 3,239 eligible children, or 41.7%). Certain regions have consistently achieved greater than 50% 3-Star visits during summer months.

Figure 3 also demonstrates the seasonality of 3-Star visits, with greater increases in the summer months. Between September 2015 and August 2016, 16,156 visits of eligible children immunized in LHDs were assessed as part of the 3-Star Report, with 52% of these visits (8,431) occurring in July and August. As 3-Star visits increased during each successive summer, the proportion of HPV-only visits correspondingly increased during the fall and winter months because children initiating the 3-dose HPV series in the summer were advised to return for dose two and three of the HPV series at recommended intervals (1–2 months later for dose 2, 6 months later for dose 3). HPV-only visit data are available in Supplementary tables.

Figure 4 shows a different perspective on the improvement in immunization practice, which is the percentage of non-HPV 1-Star visits during the same period and on the same scale as Figure 3. Here, the consistent decline in the proportion of non-HPV 1-Star visits across years and





Figure 4. Percentage of non-HPV 1-Star visits each month among VFCeligible children aged 11–13 years who received a Tdap, Men-ACWY, and/ or HPV vaccine at a Tennessee LHD, January 2013 through August 2016. Months of the year are on the *x*-axis, and percentages are on the *y*-axis.

throughout all seasons is clear. Exact numbers are provided in Supplementary tables.

No formal evaluation has been conducted of exactly how report recipients have chosen to use this monthly measurement and feedback. However, during TIP leadership site visits to LHDs across the state, staff have been asked how they are using the reports. One LHD that consistently had the lowest 3-Star rates established a temporary incentive program in February 2014. They provided the nurse with the most 3-Star visits each month a certificate of excellence and a small gift card. In that region, 3-Star visits, which ranged from 0% to 7% between July 2013 and January 2014, spiked to almost 50% in March 2014, before settling to around 20% during all summer months thereafter (results not shown). Other nursing supervisors have reported how they have tried different approaches to convincing parents and teens to receive HPV vaccine in order to increase their 3-Star rates. They also have shared anecdotally how they are pleased when they walk out of a room and know that they accomplished a 3-Star visit that will show up on the next report.

Discussion

The TIP's 3-Star Report represents a QI tool that leverages the state's existing public health informatics infrastructure, PTBMIS, to encourage LHD nurses to administer HPV and Men-ACWY in the same visit with the required Tdap vaccine. The implementation of this QI effort has corresponded with a substantial increase in 3-Star visits in LHDs across the state of Tennessee from 2013 to 2016, in particular during the summer surges of back-to-school vaccination visits for students entering seventh grade, peaking at 42% of visits in July and August of 2016. While the findings cannot be interpreted to assert causality, the observed increases in the proportion of 3-Star Visits have programmatic significance as successful performance on this QI measure.

Many states, including Tennessee, have faced challenges to achieve high vaccination rates against HPV compared to Tdap and Men-ACWY, leaving unvaccinated teens to face preventable cancer diagnoses later in life.¹⁴ Ideally, HPV vaccines are given at the age of 11-12 years, when the preteen immune system produces a more robust immune response than older immune systems and when very few have been exposed to HPV.²² As a vaccine against a virus that is often acquired through intimate skin-to-skin contact and with potential cancer diagnoses occurring later in adulthood, it may appear easier to postpone or decline for preteens than vaccines against pertussis or meningitis that may seem more immediate threats. Unfortunately, missed opportunities can result in young men and women delaying vaccination until after they acquire a serious strain of HPV or they may never be vaccinated and remain fully susceptible to HPV-related cancers.

Common strategies to improve vaccine coverage within practices that care for preteens and teens include practicelevel immunization coverage assessments, reminders for ageeligible patients, and recall notices to patients overdue for vaccinations.^{18,19} These strategies work best for clinics that serve as a preteen's medical home. In Tennessee, most LHDs do not function as a medical home, but they do offer immunizations to preteens and teens if they visit. LHDs experience a seasonal surge of preteen immunization visits, with more than half of such visits for the year taking place during the summer back-to-school season when state regulations require all students entering seventh grade to provide proof of a Tdap vaccination. Students without a medical home or whose primary healthcare provider cannot see them promptly may visit any LHD to be vaccinated.

TIP's experience with the 3-Star Report and the corresponding substantial increase in 3-Star visit rates shows how straightforward audit and feedback on a well-selected performance measure can be used to motivate continuous QI efforts in immunization. Tennessee faced substantial barriers to the use of these more comprehensive traditional measures of immunization performance. Because the state immunization information system (IIS) is a voluntary reporting system, it remains too incomplete for any type of population assessment. Because LHDs are often only an occasional source for immunization, primarily in the back-to-school rush for this age group, efforts on the part of LHD staff must be opportunistic and focused on the single encounter. To our advantage, because TIP has access to standardized data from all LHDs statewide, it is possible to generate consistent reports over time of performance at the state and local levels across all public health sites, generating a sense of friendly competition.

TIP has deliberately encouraged flexible use of the feedback reports. This report allows them to explore different strategies over time. In one case, individual nurses were identified and rewarded for their performance, a rare opportunity in clinics that do not typically examine or highlight frontline clinic nurse excellence. In other cases, individual regions emerged as high performers as a result of other concerted efforts led by LHD leadership teams. TIP encourages regions wishing to improve to learn from more successful peers. The report will continue to be refined to respond to LHD staff feedback.

To our knowledge, the 3-Star report is unique. However, the concepts behind it could be adapted to any individual clinic or network of clinics interested in increasing preteen immunization rates through bundling and interested in rewarding staff for strong recommendations for HPV vaccine. An elaborate system is not necessary to track individual immunization visits.

The 3-Star Report is not without limitations. Because this report tracks only vaccines administered during one visit, it is not possible to know whether a specific child needed fewer than three vaccines. For this reason, no target proportion of 3-Star visits was established. However, if a child comes to the LHD for a school-required Tdap, it is likely that they have not yet had any of the preteen vaccines. In addition, while the report documents HPV-only visits, it does not specifically attempt to track HPV completion rates in individual patients, given the limitations of the information available in the PTBMIS.

Furthermore, since this was a QI project and not a controlled research study, the data cannot be used to test for cause and effect or to prove that observed increases were only due to the 3-Star Report. The analysis cannot account for external factors that may have also contributed to increased parent receptiveness when the public health nurses made a 3-Star recommendation. Throughout the three years since the report was introduced, multiple efforts were ongoing to help parents understand the importance of timely HPV vaccination of boys and girls. Initiatives at the national level have been led by the Centers for Disease Control and Prevention, American Cancer Society, American Academy of Pediatrics (AAP), and other organizations. Several smaller scale efforts in focused geographic areas in the state have been led by the TIP, the Cervical Cancer Free Tennessee coalition, Vanderbilt-Ingram Cancer Center, Meharry Medical College, Cumberland Pediatric Foundation, Knox County Health Department, University of Tennessee, and others. In 2016, one of the manufacturers also introduced a new national advertising campaign directed at parents. Such educational efforts may have made it more likely that parents would accept the recommendation for on-time immunization with HPV vaccine. However, the relatively stagnant HPV vaccination coverage rates for the state suggest that there has not been a widespread cultural shift in parental acceptance during this time period that would explain the increases that were observed in LHD clinics.

HPV vaccine is capable of drastically reducing HPVrelated cancers in the next generation, but has been exceptionally underutilized. While many parents may have questions about it, we also know that most will accept it when it is recommended and used the same way as other routine

vaccines.¹⁷ A simple QI report such as this places value on the bundled visit and keeps staff focused on the expectation that the three vaccines are given to all preteens at their first immunization visit.

Conclusion

Effective strategies to increase the low rates of preteen immunization against HPV include bundling of all routinely recommended vaccines at one visit, a strong provider recommendation for immunization, and periodic audit and feedback to providers.^{13,14} The 3-Star Report demonstrates that feedback strategies do not require sophisticated and comprehensive systems to be effective. The measurement of performance at an individual visit focuses the public health nurse's attention on the manageable challenge of an individual encounter. The key to the impact of this report is its relative simplicity and consistency, as well as its documentation of trends over time and comparisons against other regions. Healthy competition can promote creative ideas that may become best practices that spread to others and increase vaccination rates broadly.

This simple and effective performance QI tool can be replicated by state immunization programs, LHDs, or primary care practices with basic electronic record systems. The principles can even be carried out on a piece of notebook paper posted in a work station. Overall, it achieves its objective without putting undue pressure on busy clinic staff. Regular performance feedback incentivizes desirable immunization provider behaviors that result in preteens receiving all recommended vaccines at a single visit by focusing on and measuring 3-Star success at each patient encounter.

Acknowledgments

The authors thank Caree McAfee for bibliographic assistance and assistance with formatting Figure 1.

Author Contributions

Conceived the report: KLM, MKF. Analyzed the data: MKF. Wrote the first draft of the manuscript: KLM. Contributed to the writing of the manuscript: KLM, MKF, PCH. Agreed with manuscript results and conclusions: KLM, MKF, PCH. Jointly developed the structure and arguments for the paper: KLM, MKF, PCH. Made critical revisions and approved the final version: KLM, MKF, PCH. All the authors reviewed and of the final manuscript.

Supplementary Material

Supplementary tables with the numerators, denominators, and percentages represented in Figures 3 and 4, as well as the numbers and proportions of HPV-only visits each month between January 2013 and August 2016 are provided.

REFERENCES

Centers for Disease Control and Prevention. *Immunization Schedules*. 2016. Available at: https://www.cdc.gov/vaccines/schedules/. Accessed October 27, 2016.



- Reagan-Steiner S, Yankey D, Jeyarajah J, et al. National, regional, state, and selected local area vaccination coverage among adolescents aged 13–17 years – United States, 2014. MMWR Morb Mortal Wkly Rep. 2015;64(29):784–92.
- Hill HA, Elam-Evans LD, Yankey D, Singleton JÅ, Kolasa M. National, state, and selected local area vaccination coverage among children aged 19–35 months – United States, 2014. MMWR Morb Mortal Wkly Rep. 2015;64(33):889–96.
- Broder KR, Cortese MM, Iskander JK, et al. Preventing tetanus, diphtheria, and pertussis among adolescents: use of tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccines recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR. 2006;55(RR-3):1–34.
- Cohn AC, MacNeil JR, Clark TA, et al. Prevention and control of meningococcal disease: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR*. 2013;62(RR-2):1–28.
- Bosch FX, Lorincz A, Muñoz N, Meijer CJ, Shah KV. The causal relation between human papillomavirus and cervical cancer. J Clin Pathol. 2002;55(4):244–65.
- Muñoz N, Castellsagué X, de González AB, Gissmann L. Chapter 1: HPV in the etiology of human cancer. *Vaccine*. 2006;24(suppl 3):1–10. doi: 10.1016/j. vaccine.2006.05.115.
- Gillison ML, Chaturvedi AK, Lowy DR. HPV prophylactic vaccines and the potential prevention of noncervical cancers in both men and women. *Cancer*. 2008;113(suppl 10):3036–46. doi: 10.1002/cncr.23764.
- 9. Petrosky E, Bocchini JA, Hariri S, et al. Use of 9-valent human papillomavirus (HPV) vaccine: updated HPV vaccination recommendations of the Advisory Committee on Immunization Practices. *MMWR*. 2015;64(11):300–4.
- Brewer NT, Fazekas KI. Predictors of HPV vaccine acceptability: a theoryinformed, systematic review. *Prev Med.* 2007;45(2–3):107–14. doi: 10.1016/j. ypmed.2007.05.013.
- Ferrer HB, Trotter C, Hickman M, Audrey S. Barriers and facilitators to HPV vaccination of young women in high-income countries: a qualitative systematic review and evidence synthesis. *BMC Public Health*. 2014;14:700. doi: 10.1186/1471-2458-14-700.
- Holman DM, Benard V, Roland KB, Watson M, Liddon N, Stokley S. Barriers to human papillomavirus vaccination among US adolescents: a systematic review of the literature. *JAMA Pediatr.* 2014;168(1):76–82. doi: 10.1001/jamapediatrics.2013.2752.

- Stokley S, Jeyarajah J, Yankey D, et al. Human papillomavirus vaccination coverage among adolescents, 2007–2013, and postlicensure vaccine safety monitoring, 2006– 2014 – United States. MMWR Morb Mortal Wkly Rep. 2014;63(29):620–4.
- President's Cancer Panel. Accelerating HPV Vaccine Uptake: Urgency for Action to Prevent Cancer. President's Cancer Panel Annual Report 2012–2013.
 2014. Available at: http://deainfo.nci.nih.gov/advisory/pcp/annualReports/ HPV/index.htm. Published 2014.
- Centers for Disease Control and Prevention. HPV: You Are the Key to Cancer Prevention. 2016. Available at: www.cdc.gov/hpv/hcp/index.html. Published 2016.
- Hull PC, Williams EA, Khabele D, Dean C, Bond B, Sanderson M. HPV vaccine use among African American girls: qualitative formative research using a participatory social marketing approach. *Gynecol Oncol.* 2014;132(suppl 1): S13-20. doi: 10.1016/j.ygyno.2014.01.046.
- Opel DJ, Heritage J, Taylor JA, et al. The architecture of provider-parent vaccine discussions at health supervision visits. *Pediatrics*. 2013;132(6):1037–1046. doi: 10.1542/peds.2013–37.
- Jacob V, Chattopadhyay SK, Hopkins DP, et al. Increasing coverage of appropriate vaccinations: a community guide systematic economic review. *Am J Prev Med.* 2016;50(6):797–808.
- Briss PA, Rodewald LE, Hinman AR, et al. Reviews of evidence regarding interventions to improve vaccination coverage in children, adolescents, and adults. The Task Force on Community Preventive Services. *Am J Prev Med.* 2000;18(suppl 1):97–140.
- Tennessee Department of Health. Results of the 2015 Immunization Status Survey of 24–Month-Old Children in Tennessee. 2016. Available at: https://www.tn.gov/assets/ entities/health/attachments/ImmunizationSurvey2015.pdf. Accessed July 22, 2016.
- Tennessee Department of Health. Childcare 12th Grade Immunization Requirements. 2016. Available at: https://www.tn.gov/health/article/childcare-12th-grade-immunization-requirements. Accessed October 27, 2016.
- Giuliano AR, Lazcano-Ponce E, Villa L, et al. Impact of baseline covariates on the immunogenicity of a quadrivalent (types 6, 11, 16, and 18) human papillomavirus virus-like-particle vaccine. *J Infect Dis.* 2007;196(8):1153–62. doi: 10.1086/521679.